

When to take SUPREP BOWEL PREP KIT

SUPREP Bowel Prep Kit is taken as a split dose (2-day) regimen. You take the first 6-ounce bottle of SUPREP the evening before your colonoscopy and the second 6-ounce bottle of SUPREP the morning of your colonoscopy. It is important to drink the additional water as recommended in the Instructions for Use. Both 6-ounce bottles are required for a complete prep. Your doctor will tell you when it is time to begin to take each dose.

How to take SUPREP BOWEL PREP KIT

- **On the evening before your procedure** (or when your doctor tells you to begin), complete steps 1 through 4 using one (1) 6-ounce bottle before going to bed
- **On the morning of your procedure**, repeat steps 1 through 4 using the other 6-ounce bottle

Both 6-ounce bottles are required for a complete preparation.

IT IS IMPORTANT TO FOLLOW ALL THE STEPS BELOW COMPLETELY.

Step 1:



Pour **ONE** (1) 6-ounce bottle of SUPREP liquid into the mixing container.

Step 2:



Add cool drinking water to the 16-ounce line on the container and mix.

NOTE: Be sure to dilute SUPREP as shown at left before you drink it.

Step 3:



Drink **ALL** the liquid in the container.

Step 4:



You **must** drink two (2) more 16-ounce containers of water over the next 1 hour.

NOTE: You **must** finish drinking the final glass of water at least 2 hours, or as directed, before your procedure.